

Seyfert 2 Galaxies with Spectropolarimetric Observations

Qiusheng Gu and Jiehao Huang

Department of Astronomy, Nanjing University, Nanjing 210093, P. R. China

qsgu, jhh@nju.edu.cn

ABSTRACT

We present a compilation of radio, infrared, optical and hard X-ray (2-10keV) data for a sample of 90 Seyfert 2 galaxies (Sy2s) with spectropolarimetric observations (41 Sy2s with detection of polarized broad lines (PBL) and 49 without PBL). Compared to Sy2s without PBL, Sy2s with PBL tend to be earlier-type spirals, and show warmer mid-infrared color and significant excess of emissions (including the hard X-ray (2-10keV), [O III] λ 5007, infrared (25 μ m) and radio). Our analyses indicate that the majority of Sy2s without PBL are those sources having less powerful AGN activities, most likely caused by low accretion rate. It implies that the detectability of the polarized broad emission lines in Sy2s may depend on their central AGN activities in most cases. Based on the available data, we find no compelling evidence for the presence of two types of Sy2s, one type of them has been proposed to be intrinsically different from Sy2s claimed in Unification Model.

Subject headings: galaxies: active — galaxies: Seyfert — galaxies: statistics

1. Introduction

In the scheme of the standard unification model, Seyfert 1 and 2 galaxies (Sy1s and Sy2s hereafter) are intrinsically the same objects and the absence of broad emission lines in Sy2s is ascribed to the obscuration by a pc-scale dusty torus oriented along the line of sight (see the reviews by Antonucci 1993). The observational evidence for this model includes the detection of polarized broad emission lines in some Seyfert 2 galaxies (Antonucci & Miller 1985; Tran 1995 and 2001; Young et al. 1996; Heisler et al. 1997; and Moran et al. 2000), the detection of broad lines in the infrared spectra of some Sy2s (Rix et al. 1990; Ruiz et al. 1994; and Veilleux et al. 1997) and the detection of a prominent photoelectric cutoff in the X-ray spectra of Sy2s indicating the presence of large columns of gas along the line of sight (Koyama et al. 1989; Awaki et al. 1991; Maiolino et al. 1998; Risaliti et

al. 1999). However, recent investigations suggest that this strictest version of unification model needs modifications. Among them, we may find, for example, the outflowing wind model (Elvis 2000), and the existence of two intrinsically different populations of Sy2s – the hidden Sy1 and the "real" Sy2 with a weak or absent Sy1 nucleus – (Tran 2001) based on a spectropolarimetric survey of the CfA and $12\mu\text{m}$ samples of Seyfert 2 galaxies. Nevertheless, Antonucci (2001) strongly argued that the evidence claimed by Tran is quite uncertain.

With the improvement of the techniques and instruments for the spectropolarimetry, now people have observed a large sample of Sy2s with less bias (e.g. Heisler et al. 1997; Moran et al. 2000). For the present, polarized broad lines (PBL) have been detected in several dozens of Sy2s, while not detected in other several dozens. Although such surveys are probably biased inherently since pre-selection was done according to the broad-band polarization, they still provide a largest sample for more meaningful analysis than before so that we may, or may not, find some compelling evidence for the proposed modifications, especially the presence of two types of Sy2s. That is what we would present in this paper.

In this work, we provide the multiwavelength data for a sample of 90 Seyfert 2 galaxies collected from recent literatures in §2, and compare the properties of host galaxies, infrared, radio and hard X-ray continua and [O III] emission in §3. The implications and discussions on the results are given in §4, and conclusions in §5 .

2. The Sample

We collect all Seyfert 2 galaxies with available spectropolarimetric data from recent literature (from 1985 to 2002), where 41 Sy2s with PBL and 49 without the detection of PBL are found. Just as most, if not all, Seyfert samples, the current spectropolarimetry sample of Sy2s (SPSS hereafter) is heterogeneous and cannot be considered complete, though it covers a wide range of AGN luminosities (about 4 orders of magnitudes).

The infrared and radio data for these SPSS sources are listed in Table 1, which contains galaxy name (column 1); the IRAS color f_{60}/f_{25} (column 2); the $25\mu\text{m}$ luminosity (column 3); the far-infrared ($40 - 120\mu\text{m}$) luminosity (column 4), determined by (Lonsdale et al. 1985)

$$L_{\text{FIR}} = 3.75 \cdot 10^5 \cdot D^2 (2.58f_{60} + f_{100})$$

where D is the distance in Mpc (the Hubble constant is taken as $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$), f_{60} and f_{100} are the flux densities at $60\mu\text{m}$ and $100\mu\text{m}$ in Jy; the radio powers at 1.49 GHz (column 5) from the NRAO/VLA Sky Survey (NVSS) (Condon et al. 1998); the mean revised morphological types (column 6) from the Third Reference Catalogue of Bright Galaxies (RC3,

de Vaucouleurs et al. 1991); and the information for PBL (columns 7 and 8). Following Lumsden et al. (2001), we adopt the infrared fluxes from BGS (Soifer et al. 1989; Sanders et al. 1995) for Sy2s with $f_{60} > 5\text{Jy}$; from Strauss et al. (1990) for those with $f_{60} < 5\text{Jy}$, and IRAS Faint Source Catalog (FSC, Moshir et al. 1992) for all other ones.

In Table 2, we present the optical and hard X-ray information as the following: column 1, the galaxy name; column 2, the blue luminosity computed with the formula (Pogge & Eskridge 1993)

$$\log L_B = 12.208 - 0.4B_T^0 + \log(1+z) + 2\log D$$

where z is the redshift of the galaxy; column 3, the luminosity of H_β emission; column 4, the luminosity of extinction-corrected $[\text{O III}]\lambda 5007$ emission given as $L_{[\text{O III}]} = 4\pi D^2 F_{[\text{O III}]}^{\text{cor}}$, where $F_{[\text{O III}]}^{\text{cor}}$ is the extinction-corrected flux of $[\text{O III}]\lambda 5007$ emission derived from the relation (Bassani et al. 1999)

$$F_{[\text{O III}]}^{\text{cor}} = F_{[\text{O III}]}^{\text{obs}} \left(\frac{(H_\alpha/H_\beta)_{\text{obs}}}{(H_\alpha/H_\beta)_0} \right)^{2.94}$$

we assume an intrinsic Balmer decrement $(H_\alpha/H_\beta)_0 = 3.0$; column 5, the references for H_β and $[\text{O III}]\lambda 5007$; column 6, gaseous absorbing column density (N_H); column 7, the absorption-corrected hard X-ray (2-10keV) luminosity for Compton-thin Seyfert 2 galaxies; column 8, the corresponding references; and column 9, Eddington luminosity acquired by $L_{\text{Edd}} = 1.51 \cdot 10^{38} M_{\text{BH}}/M_\odot \text{ ergs}^{-1}$. The central black hole masses are estimated by use of the $M_{\text{BH}} - \sigma$ relation by Merritt & Ferrarese (2000), which is

$$M_{\text{BH}} = 1.3 \times 10^8 \left(\frac{\sigma}{200 \text{ km s}^{-1}} \right)^{4.72} M_\odot$$

Nuclear velocity dispersions (σ) are taken from Nelson & Whittle (1995) and McElroy (1995) if available.

3. The results

3.1. Properties of host galaxies

Following the approach taken by Storchi-Bergmann et al (2001) to evaluate the morphological type distribution for their Seyfert sample, we display the histogram distributions of morphological types for our Sy2s *with* and *without* PBL (SPSS1 and SPSS0 for simplicity hereafter) in Fig 1a and 1b, respectively. The statistical median of their morphological type distributions are Sa and Sab for SPSS1 and SPSS0, showing with stars in Fig1a and 1b, respectively. The SPSS1 (Sy2s *with* PBL) obviously tend to be earlier-type spirals.

In Fig.2, we present the cumulative distributions of blue luminosities (in unit of $L_{\odot}$) for SPSS1 and SPSS0, denoted with filled and open circles, respectively. The two-sample Kolmogorov-Smirnov (KS) test, *kolmov* task in IRAF¹, shows that there exists no significant difference between the two subsamples. The same conclusion could be reached with the Kaplan-Meier estimator, *kmestimate* task in IRAF, yielding L_B of 10.475 ± 0.066 and 10.349 ± 0.085 for SPSS1 and SPSS0, respectively. The basic statistical results presented in this paper are summarized in Table 3.

We also compare the column densities (N_H) for SPSS1 and SPSS0. Due to the existence of censored data in column densities (20 sources with lower limits and 1 source with upper limit, NGC 7590), the survival analysis methods (ASURV Rev 1.2², Isobe & Feigelson, 1990) has been used for the statistical analysis. A possibility of 79% for the test is given by the Gehan's Generalized Wilcoxon Test - Hypergeometric Variance (GGW test, one kind of *asurv* tests), suggesting little difference in $\log N_H$ distributions between SPSS1 and SPSS0. Following Alexander (2001) and Tran (2001), we may also estimate the mean $\log N_H$ of $23.63 \pm 0.20 \text{ cm}^{-2}$ and $23.56 \pm 0.35 \text{ cm}^{-2}$ ($23.67 \pm 0.35 \text{ cm}^{-2}$ if NGC 7590 is excluded, since *asurv* could not deal with the case which contains both upper and lower limits) for the two subsamples, respectively, which leads to the same conclusion on the column density (N_H). Indeed, Alexander (2001), Gu et al. (2001) and Tran (2001) have derived the similar results for samples with smaller size.

3.2. Indicators of AGN activities

During the last two decades, there have been several attempts to determine which emissions are truly isotropic for Seyfert galaxies. Among various emissions at different wavelength bands, the [O III] λ 5007, infrared and hard X-ray (2-10keV) continua have been found (Dahari & De Robertis 1988, Keel et al. 1994, Mulchaey et al. 1994, Alonso-Herrero et al. 1997) to indicate similar distributions for Sy1s and Sy2s, implying their isotropic properties so as to be good indicators of the intrinsic nuclear luminosity.

Recent investigation have found that about (30-50)% of Seyfert 2 galaxies show nuclear starburst activities (Storchi-Bergmann et al. 2000; Gonzalez Delgado et al. 2001). It follows that the infrared radiation may not be a good indicator for AGN luminosity, because of

¹IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

²<http://www.astro.psu.edu/statcodes/asurv>

the contamination from nuclear star-forming activities. On the other hand, the incomplete isotropy of [O III] emission caused by obscuration from host galaxies brings about a suggestion (Maiolino et al. 1998) to take the extinction-corrected [O III] luminosity as an indicator of the nuclear activity.

The same is true for the hard X-ray (2-10keV) continua, in a sense that they are heavily absorbed. About half of Sy2s in the Local Universe are Compton-thick with absorbing column densities $N_H > 10^{24} \text{ cm}^{-2}$ (Maiolino et al. 1998; Bassani et al. 1999; Risaliti, Maiolino & Salvati 1999). In the case of Compton-thick Seyfert 2 galaxies, the direct hard X-ray continua are completely absorbed, implying that this indicator of AGN activities works well for Compton-thin sources only. What is more, the effect of absorption on X-ray emission should be corrected, such as deriving the extinction-corrected [O III] mentioned above.

3.3. Statistics on hard X-ray – radio emissions

Following the above considerations, we perform statistics on various indicators of AGN activities for SPSS1 and SPSS0 subsamples. The results are illustrated in Fig 3 - Fig 6, where one could find cumulative distributions of extinction-corrected [O III] luminosities (in Fig 3), mid-infrared flux ratios (f_{60}/f_{25}), 25 μm luminosities and radio powers at 1.49 GHz (in Fig 4a, b and c), respectively. It is evident from the comparisons that SPSS1 differ from SPSS0 dramatically. The Sy2s *with* PBL (SPSS1) show more powerful [O III] emission, much hotter mid-infrared color and significant excesses of 25 μm and radio emissions. Indeed, the KS test or GGW test shows that the SPSS1 and SPSS0 subsamples are drawn from the same parent population with a possibility $P_{\text{null}} < 0.0001$. The statistical results are given in Table 3.

In Fig 5, we show a plot of the absorption-corrected 2-10 keV luminosity vs extinction-corrected [O III] luminosity for Compton-thin Seyfert 2 galaxies in SPSS1 (in filled circles) and SPSS0 (in open circles) subsamples. It is instructive to find that Compton-thin Sy2s *with* and *without* PBL are located in separate regions. The mean $\log L_{2-10\text{keV}}$ for SPSS1 and SPSS0 sources are $43.0 \pm 0.14 \text{ erg s}^{-1}$ and $41.22 \pm 0.32 \text{ erg s}^{-1}$, respectively, clearly indicating that the nuclear activities in SPSS1 are much more powerful than that in SPSS0. The GGW test of $P_{\text{null}} < 0.0001$ provides full support to this claim.

Probably related to the above finding is what we show in Fig 6, a plot of extinction-corrected [O III] emission against Eddington luminosity (L_{Edd}) for SPSS1 and SPSS0 subsamples. In accordance with the strong tendency manifested in Fig 5, what Fig 6 illustrates is that the majority of SPSS1 sources have higher $L_{[\text{OIII}]} / L_{\text{Edd}}$. That is, 11 out of 14 SPSS1

sources have $L_{[\text{OIII}]} / L_{\text{Edd}} > 10^{-4}$, while 12 Of 17 SPSS0 $< 10^{-4}$. It suggests that the accretion rates in SPSS1 sources might be higher than that in SPSS0 on average.

4. Discussions

It has long been known that $L_{\text{FIR}} / L_{\text{B}}$ is a good star formation indicator for spiral galaxies (see, e.g. Gu et al. 1999; Lei et al. 2000). We find that the distributions of $L_{\text{FIR}} / L_{\text{B}}$ for SPSS1 and SPSS0 are similar, with the mean $L_{\text{FIR}} / L_{\text{B}}$ of -0.275 and -0.298, respectively. It implies that the detectability of PBL does not directly relate to the star-forming activities in host galaxies. In fact, we may list both SPSS1 and SPSS0 sources showing nuclear starburst activities, e.g. Mrk 477, Mrk 533, IC 3639 in SPSS1, and NGC 5135, NGC 7130 in SPSS0 (Gonzalez Delgado et al. 2001).

The detectability of a hidden BLR in Sy2s has been suggested (Heisler et al. 1997) having relation to the inclination of the torus determined by the $60 \mu\text{m}$ to $25 \mu\text{m}$ flux ratio. However, by comparing the column densities inferred from hard X-rays, Alexander (2001) argued against the direct relation of the detectability of the PBL to IR colors and claimed that the contribution from host galaxies would both make the IR color cooler and dilute the nuclear activity. In the mean time, Gu, Maiolino & Dultzin-Hacyan (2001) find that the detectability of PBL depends on the relative contribution/dilution from host galaxies. Lumsden et al. (2001) also suggested that the detectability of PBL is largely due to a combination of AGN activity, the obscuring density and the relative contribution of host galaxies.

By analyzing various indicators of AGN activities, we find in this work that mid-infrared, radio, hard X-ray, and [O III] luminosities of SPSS1 are significantly larger than SPSS0, indicating obviously that the detectability of PBL is most likely due to the AGN activities. Indeed, the mean absorption-corrected X-ray luminosity in 2-10keV energy band for SPSS1, $L_{2-10\text{keV}}^c \sim 10^{43} \text{ erg s}^{-1}$, is the typical X-ray luminosity for Seyfert 1 galaxies (Nandra et al., 1997). The mean $L_{2-10\text{keV}}^c$ of SPSS0, on the contrary, is nearly 2 order of magnitudes smaller, most of them show very low accretion rates as Fig 6 illustrated. What we can infer from these analyses is that the majority of SPSS0 (Sy2s *without* PBL) sources are probably low-luminosity AGNs (LLAGNs). The more powerful the AGNs are, the easier it is to detect PBL, which is also consistent with our finding of the morphological difference of Sa from Sab between SPSS1 and SPSS0. Galaxies with earlier Hubble types have mirrors that are easier to be detected.

Our sample is an amalgamation of different observations with diverse quality of spec-

tropolarimetric data, varying from object to object determined by the brightness, observers, integration time, and a host of other factors. For example, Ruiz et al. (1994) have found no evidence for broad H_α and H_β components in the polarized light for Mrk 334, yielding a maximum of $\sim 0.6\%$ of the light in H_α as polarized broad-line flux. For other Sy2s with detected polarized broad lines, the mean percentage of 1% - 5% has been reported. Needless to say that it is necessary to evaluate the sensitivities to detect polarized broad lines in different groups from which our data are taken. The difficult situation we are faced with is that only one group (Young et al 1996) has presented detailed information for their polarized broad emission components, so that we can derive their sensitivity – the detected minimum polarized broad H_α flux is $1.2 \cdot 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$ for NGC 7674 and their mean percentage of polarized broad H_α emission to total H_α emission is $4.7 \pm 0.8 \%$. For three groups (Moran et al 2000, Moran et al 2001, Tran 2001, Lumsden et al 2001) reporting no detailed polarized properties, to compare the common sources that they observed would be an adequate approach to evaluate their sensitivities. The analyzed results are summarized in Table 4.

At first glance, one could find that there are 16 common Sy2s in both Tran’s and Lumsden’s samples and 13 in both Tran’s and Moran’s samples, and that detectabilities of PBL for these common sources are the same, which indicates comparable sensitivities of these three samples. On the other hand, 3 Sy2s (IRAS 00521-7054, IRAS 04385-0828 and NGC 5506) with PBL reported by Tran (2001) are not listed in Young’s sample as Sy2s *with* PBL. What is more, two Sy2s (NGC 5347 and NGC 5929) listed as ”non-HBLR” Sy2s in Tran’s sample have been discovered (Moran et al. 2001) showing faint polarized broad emission lines with high S/N data.

The conclusions we could reach from the above analyses is that the sensitivity of Moran’s sample (2001) is higher than that of Tran’s (2001), both of which are higher than that of Young’s sample (1996). Since the polarized fluxes of broad emission lines represent a certain fraction of the total flux, typically a few percents. it should be true to say that the Sy2s *without* detection of PBL would be preferentially harder to be detected for a given sensitivity. In other words, for less luminous Sy2s without detection of PBL, one would need to reach sensitivities that would allow us to detect the *same percentage polarization*. Therefore, to attribute those Sy2s without detection of PBL to non-hidden BLR in those sources would be premature. What we can point out clearly is that *we have found no compelling evidence for the presence of two types of Sy2s as claimed by Tran(2001)*.

Antonucci (2001) has repeatedly made the point that, to be compelling, evaluations of one sample vs. another must be made for samples that are selected by a property which is thought to be isotropic. Due to the limitation of our sample selection (heterogeneous

with different spectropolarimetric sensitivities), it is impossible to interpret our statistical results in the configuration of testing the unified model for AGN. On the contrary, the conclusions we reached above force us to claim at the present time that *we have found no compelling evidence to argue for modifications to the unified model of AGN*. The non-detection of PBL is probably due to the low AGN activities. Based on the analyses presented in this paper, we can not rule out the possibility of the combination of the weakness of AGN and commensurate obscuration/dilution from host galaxies (Lumsden & Alexander 2001, Gu Maiolino & Dultzin-Hacyan 2001).

5. Conclusions

In this paper, we collect radio, infrared, optical and hard X-ray data for a sample of 90 Seyfert 2 galaxies with spectropolarimetric observations. Out of these 90 objects, 41 show polarized broad lines (most likely ascribed to scattering of the broad line region) and 49 do not. Compared to Sy2s without PBL, Sy2s with PBL tend to be earlier-type spirals, and show warmer mid-infrared color and significant excess of emissions (including the hard X-ray(2-10keV), [O III] λ 5007, infrared (25 μ m) and radio), while their distributions of blue luminosity and absorbing column density are similar. Our analyses suggest that the majority of Sy2s without PBL are those sources having less powerful AGN activities, most likely caused by low accretion rate. It implies that the detectability of the polarized broad emission lines in Sy2s may depend on their central AGN activities in most cases. Based on the available data, we find no compelling evidence for the presence of two types of Sy2s, one type of them has been proposed to be intrinsically different from Sy2s claimed in Unification Model.

The authors are very grateful to the anonymous referee for his/her critical comments and instructive suggestions, which significantly strengthened the analyses in this paper. We also thank Luis Ho, Xiaoyang Xia and Zupan Deng for their valuable discussion and suggestion. This work has been supported by the National Natural Science Foundation of China under grant 10103001 and the National Key Basic Research Science Foundation (NKBRF G19990754). This research has made use of NASA’s Astrophysics Data System Abstract Service and the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Acker A., Stenholm B. & Veron P., 1991, A&AS, 87, 499.
- Alexander D.M., 2001, MNRAS, 320, L15.
- Almudena Prieto M., Perez Garcia A.M. & Rodriguez Espinosa J.M., 2002, MNRAS, 329, 309.
- Alonso-Herrero A., Ward M. J., Kotilainen J.K., 1997, MNRAS, 288, 977.
- Antonucci R. 1993, ARA&A, 31, 473.
- Antonucci R. & Miller J.S., 1985, ApJ, 297, 621.
- Antonucci R., 2001, astro-ph/0110343.
- Awaki H., Koyama K., Inoue H. & Halpern J.P., 1991, PASJ, 43, 195.
- Awaki H., Ueno S., Taniguchi Y. & Weaver K., 2000, ApJ, 542, 175.
- Bassani L., Dadina M., Maiolino R., et al., 1999, ApJS, 121, 473.
- Collinge M.J. & Brandt W.N., 2000, MNRAS, 317, L35.
- Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B. & Broderick J. J., 1998, AJ, 115, 1693.
- Cruz-Gonzalez I., Carrasco L., Serrano A., et al., 1994, ApJS, 94, 47.
- Dahari O., 1985, ApJS, 57, 643.
- Dahari O. & De Robertis M. M., 1988, ApJS, 67, 249.
- De Grijp M. H. K., Keel W.C., Miley G.K., & Lub J., 1992, A&AS, 96, 389.
- de Vaucouleurs G., de Vaucouleurs A., Corwin J. R., Buta R. J., Paturel G. & Fouque P., 1991, Third Reference Catalogue of Bright Galaxies, 1991, New York : Springer-Verlag. (RC3)
- Elvis M., 2000, ApJ, 545, 63.
- Gonzalez Delgado R., Heckman T. & Leitherer C., 2001, ApJ, 546, 845.
- Gu Q., Huang J. & Ji L., 1999, Ap&SS, 260, 389.

- Gu Q., Maiolino R. & Dultzin-Hacyan D., 2001, *A&A*, 366, 765.
- Heisler C.A., Lumsden S.L. & Bailey J.A., 1997, *Nature*, 385, 700.
- Ho L., Filippenko V. & Sargent W. L. W., 1997, *ApJS*, 112, 315.
- Inglis M.D., Brindle C., Hough J.H., Young S., et al., 1993, *MNRAS*, 263, 895.
- Isobe T. & Feigelson E.D., 1990, *BAAS*, 22, 917.
- Kay L. E. & Moran E.C., 1998, *PASP*, 110, 1003.
- Keel W.C., de Grijp M.H.K., Miley G.K. & Zheng W., 1994, *A&A*, 283, 791.
- Koyama K., Inoue H., Tanaka Y., Awaki H., Takano S., Ohashi T. & Matsuoka M., 1989, *PASJ*, 41, 731.
- Lei S., Huang J., Zheng W., Ji L. & Gu Q., 2000, *ApJ*, 544, L31.
- Levenson N. A., Weaver K. A. & Heckman T. M., 2001, *ApJ*, 550, 230.
- Lonsdale C.J., Helou G., Good J.C. & Rice W.L., 1985, *Catalogued Galaxies and Quasars Observed in the IRAS Survey*, Washington D.C.
- Lonsdale C.J., 1992, *ApJ*, 391, 629.
- Lumsden S. L., Heisler C. A., Bailey J. A., Hough J. H., & Young S., 2001, *MNRAS*, 327, 459.
- Lumsden S. L. & Alexander D. M., 2001, *MNRAS*, 328, L32.
- Merritt D. & Ferrarese L., 2000, *ApJ*, 547, 140.
- Maiolino R., Ruiz M, Rieke G.H. & Keller L.D., 1995, *ApJ*, 446, 561.
- Maiolino R., Salvati M., Bassani L., et al., 1998, *A&A*, 338, 781.
- McElroy D. B., 1995, *ApJS*, 100, 105.
- Miller J.S. & Goodrich R.W., 1990, *ApJ*, 355, 456.
- Moran E.C., Halpern J.P., Bothun G.D. & Becker R.H., 1992, *AJ*, 104, 990.
- Moran E.C., Barth A.J., Kay L.E. & Filippenko A.V., 2000, *ApJ*, 540, L73.
- Moran E.C., Kay L.E., Davis M., Filippenko A.V. & Barth A., 2001, *ApJ*, 556, L75.

- Moshir M., Kopman G. & Conrow T.A.O., 1992, IRAS Faint Source Survey, Explanatory supplement version 2, JPL D10015 8/92, JPL, Pasadena.
- Mulchaey J., Koratkar A., Ward M., Wilson A., et al., 1994, ApJ, 436, 586.
- Nandra K., George I.M., Mushotzky R., Turner T.J. & Yaqoob T., 1997, ApJ, 476, 70.
- Nelson C. H. & Whittle M., 1995, ApJS, 99, 67.
- Oliva E., Salvati M., Moorwood A.F.M. & Marconi A., 1994, A&A, 288, 457.
- Polletta M., Bassani L., Malaguti G., Palumbo G.G.C. & Caroli E., 1996, ApJS, 106, 399.
- Pogge R.W. & Eskridge P.B., 1993, AJ, 106, 1405.
- Risaliti G., Maiolino R. & Salvati M., 1999, ApJ, 522, 157.
- Risaliti G., Gilli R., Maiolino R. & Salvati M., 2000, A&A, 357, 13.
- Rix H., Rieke G., Rieke M. & Carleton N.P., 1990, ApJ, 363, 480.
- Ruiz M., Rieke G. H. & Schmidt G. D., 1994, ApJ, 423, 608.
- Sanders D.B., Egami E., Lipari S., Mirabel I., Soifer B.T., 1995, AJ, 110, 1993.
- Soifer B.T., Bochmer L. Neugebauer G., Sanders D.B., 1989, AJ, 98, 766.
- Storchi-Bergmann T. & Griselda Pastoriza M., 1989, ApJ, 347, 195.
- Storchi-Bergmann T., Kinney A.L. & Challis P., 1995, ApJS, 98, 103.
- Storchi-Bergmann T., Raimann D., Bica E.L.D. & Fraquelli H.A., 2000, ApJ, 544, 747.
- Storchi-Bergmann, T, Gonzalez Delgado, R. M., Schmitt, H. R., Cid Fernandes, R., & Heckman, T. 2001, ApJ, 559, 147
- Strauss M.A., Davis M., Yahil A. & Huchra J.P., 1990, ApJ, 361, 49.
- Terlevich R., Melnick J., Masegosa J., Moles M. & Copetti M., 1991, A&AS, 91, 285.
- Tran H.D., Miller J.S. & Kay L.E., 1992, ApJ, 397, 452.
- Tran H.D., 1995, ApJ, 440, 565.
- Tran H.D., Brotherton M.S., Stanford S.A., et al., 1999, ApJ, 516, 85.
- Tran H.D., 2001, ApJ, 554, L19.

- Vaceli M.S., Viegas S.M., Gruenwald R. & Souza R.E., 1997, *AJ*, 114, 1345.
- Vanzi L., Bagnulo S., Floch E., Maiolino R., et al., 2002, *A&A*, in press (astro-ph/0202435).
- Veilleux S., Kim D.C., Sanders D., Mazzarella J.M. & Soifer B.T., 1995, *ApJS*, 98, 171.
- Veilleux S., Goodrich R.W. & Hill G.J., 1997, *ApJ*, 477, 631.
- Véron-Cetty M.P. & Véron P., 2001, *Quasars and Active Galactic Nuclei* (10th Ed.), *A&A*, 374, 92.
- Whittle M. 1992, *ApJS*, 79, 49.
- Young S., Hough J.H., Efstathiou A., Wills B.J., Bailey J.A., Ward M.J. & Axon D.J., 1996, *MNRAS*, 281, 1206.




Fig. 1.— The histogram distribution of the morphological types for Sy2s with PBL (a) and for Sy2s without PBL (b).




Fig. 2.— The cumulative distribution of the blue luminosities for Sy2s with PBL (filled circles) and for Sy2s without PBL (open circles).




Fig. 3.— The cumulative distribution of luminosities of extinction-corrected [O III] λ 5007 emission. Symbols have the same coding as in Fig 2.




Fig. 4.— The cumulative distribution of infrared ratio f_{60}/f_{25} (a), mid-infrared ($25 \mu\text{m}$) luminosity (b) and radio 1.49 GHz continuum power (c). Symbols have the same coding as in Fig 2.




Fig. 5.— The distribution of absorption-corrected 2-10keV luminosities against extinction-corrected [O III] luminosities for Compton-thin Seyfert 2 galaxies. Symbols have the same coding as in Fig 2.




Fig. 6.— The distribution of Eddington luminosities against [O III] luminosities. Symbols have the same coding as in Fig 2.

Table 1. Infrared and radio data for Seyfert 2 galaxies

Name (1)	f_{60}/f_{25} (2)	$L_{25\mu\text{m}}^a$ (3)	L_{FIR}^a (4)	$L_{1.49\text{GHz}}^b$ (5)	Type (6)	PBL? (7)	References (8)
Mrk 334	4.038	10.505	10.674	29.434	pec	no	R94
NGC 34	6.756	10.751	11.143	29.705	Sc	no	H97
IRAS 00198-7926	2.488	11.603	11.547		pec	no	T01
Mrk 348	1.419	10.161	9.855	30.132	SA(s)0/a	yes	M90
IRAS 00521-7054	1.270	11.357	11.064		E-S0	yes	T01
NGC 424	1.183	10.122	9.740	28.752	(R)SB(r)0/a	yes	M00
NGC 513	6.326	9.856	10.348	29.621	Sbc	yes	T01
NGC 591	4.327	9.827	10.094	29.187	(R')SB0/a	yes	M00
Mrk 573	1.593	10.150	9.906	29.133	(R)SAB(rs)0	no	T01
IRAS 01475-0740	1.095	10.252	9.798	30.278	E-S0	yes	T01
NGC 788	0.992	9.765	9.336		SA(s)0/a	yes	K98
NGC 1068	2.160	10.865	10.789	30.130	(R)SA(rs)b	yes	A85
NGC 1144	8.724	10.459	11.087	30.391	S pec	no	T01
Mrk 1066	4.857	10.321	10.578	29.467	(R)SB(s)0	no	M00
IRAS 02581-1136	1.067	10.407	10.168	29.175	SAB(rs)a	yes	T01
NGC 1241	8.294	9.767	10.381		SB(rs)b	no	T01
NGC 1320	2.189	9.722	9.644	27.974	Sa	no	T01
NGC 1358	3.048	9.115	9.302	28.820	SAB(r)0/a	no	M00
NGC 1386	4.027	8.729	8.962	27.637	SB(s)0	no	M00
IRAS 03362-1642	2.133	10.606	10.588	29.370		no	T01
IRAS 04103-2838	3.388	11.646	11.724	30.539		no	Y96
IRAS 04210+0400	2.378	10.487	10.697	30.295		no	Y96
IRAS 04229-2528	3.687	10.478	10.633	29.547		no	Y96
IRAS 04259-0440	2.798	10.311	10.297		E-S0	no	H97
IRAS 04385-0828	1.766	10.357	10.125	28.882	S0	yes	T01
NGC 1667	8.914	9.966	10.636	29.506	SAB(r)c	no	M00
NGC 1685	4.477	9.456	9.726	28.753	SB(r)0/a	no	M00
IRAS 05189-2524	3.986	11.578	11.721	29.999	pec	yes	H97
Mrk 3	1.300	10.515	10.171	30.600	S0	yes	M90
NGC 2273	4.881	9.631	9.932	28.803	SB(r)a	yes	M00
ESO 428-G014	2.486	9.376	9.370	28.542	SA0	no	M00
Mrk 1210	0.984	10.283	9.792	29.571	Sa	yes	T92
IRAS 08277-0242	3.449	10.620	10.742	30.028		no	Y96
NGC 3081				27.731	(R1)SAB(r)0/a	yes	M00
NGC 3079	14.014	9.521	10.336	29.359	SB(s)c	no	T01
NGC 3281	2.645	10.257	10.250	29.248	SAB(rs+)a	no	M00
IRAS 10340+0609	1.556	9.309	9.240			no	M00
NGC 3362				29.376	SABc	no	T01
UGC 6100	2.842	10.012	10.182	29.265	Sa	no	T01
IRAS 11058-1131	2.387	10.763	10.700	29.592		yes	V01
NGC 3660	7.780	9.315	9.905	28.513	SB(r)bc	no	T01
NGC 3982	8.258	8.918	9.531	28.226	SAB(r)b	no	M00
NGC 4117				27.020	S0	no	M00
Was 49b	1.446	10.856	10.617	30.755		yes	T92
NGC 4388	2.939	10.139	10.244	29.188	SA(s)b	yes	H97
NGC 4501	5.953	9.990	10.559	29.468	SA(rs)b	no	T01
NGC 4507	3.953	9.952	10.168	29.190	SAB(s)ab	yes	M00
IC 3639	2.953	10.222	10.308	29.265	SB(rs)bc	yes	H97

Table 1—Continued

Name (1)	f_{60}/f_{25} (2)	$L_{25\mu\text{m}}^{\text{a}}$ (3)	$L_{\text{FIR}}^{\text{a}}$ (4)	$L_{1.49\text{GHz}}^{\text{b}}$ (5)	Type (6)	PBL? (7)	References (8)
NGC 4941	4.098	8.480	8.812	27.629	(R)SAB(r)ab	no	M00
MCG -3-34-64	2.058	10.683	10.549	30.158	SB?	yes	T01
NGC 5135	6.732	10.422	10.886	29.824	SB(l)ab	no	H97
NGC 5194	6.587	9.480	10.035	28.441	SA(s)bc	no	T01
NGC 5252				29.212	L	yes	Y96
NGC 5256	6.363	10.729	11.138	30.283	pec	no	T01
NGC 5283				28.355	S0	no	M00
Mrk 1361	3.905	10.406	10.569	29.297	SB	no	H97
IRAS 13452-4155	2.272	10.857	10.735			no	Y96
NGC 5347	1.253	9.656	9.442	27.852	(R')SB(rs)ab	yes	M01
Mrk 463E	1.514	11.356	11.087	31.272		yes	M90
Circinus	3.640	9.513	9.662		SA(s)b	yes	V01
NGC 5506	2.014	9.947	9.809	29.360	Sa	yes	T01
NGC 5643	5.025	9.552	9.895		SAB(rs)c	no	M00
NGC 5695	4.388	9.203	9.606	28.397	SBb	no	M00
Mrk 477	2.507	10.674	10.672	30.230		yes	T92
NGC 5728	10.195	9.636	10.275	29.047	(R1)SAB(r)a	no	M00
ESO 273-IG04	2.767	11.172	11.174			yes	V01
NGC 5929	5.642	9.889	10.253	29.220	Sab	yes	M01
NGC 5995	2.824	10.724	10.808	29.561	S(B)c	yes	T01
IRAS 15480-0344	1.468	10.601	10.574	29.863	S0	yes	T01
IRAS 17345+1124	2.469	11.487	11.855	31.782		yes	V01
NGC 6552	3.644	10.342	10.491	29.640	SB?	yes	T01
IRAS 19254-7245	3.955	11.484	11.667			no	H97
NGC 6890	4.923	9.480	9.873		(R')SA(r)ab	no	M00
IRAS 20210+1121	2.421	11.424	11.337	30.485		no	Y96
IRAS 20460+1925	1.672	12.014	11.865	31.064		yes	Y96
IC 5063	1.528	10.468	10.170		SA(s)0+	yes	I93
NGC 7130	7.757	10.532	11.038	29.977	Sa	no	H97
NGC 7172	6.372	9.632	10.100	28.731	Sa	no	H97
IRAS 22017+0319	1.607	11.206	10.953	30.060		yes	T01
NGC 7212	4.083	10.489	10.736	30.188		yes	T92
MCG -3-58-7	2.774	10.718	10.773	29.374	(R')SAB(s)0/a	yes	T01
IRAS 23060+0505	2.700	11.881	11.833	30.584	pec	yes	V01
NGC 7496	5.536	9.531	9.890		(R':)SB(rs)bc	no	H97
IRAS 23128-5919	6.792	11.276	11.666			no	Y96
IC 5298	4.654	10.943	11.205	29.715		no	L01
NGC 7582	6.895	10.078	10.531		(R'1)SB(s)ab	no	H97
NGC 7590	9.065	9.182	9.844		S(r?)bc	no	H97
NGC 7672	2.961	9.271	9.646	28.382	Sb	no	M00
NGC 7674	2.809	10.996	11.057	30.572	SA(r)bc	yes	M90
NGC 7682	2.034	9.725	9.678	29.545	SB(r)ab	yes	T01

^aLuminosities in units of $L_{\odot}$.

^bRadio 1.49GHz luminosity in unit of $\text{erg s}^{-1} \text{Hz}^{-1}$

Note. — REFERENCES FOR PBL: A85 = Antonucci & Miller (1985); H97 = Heisler et al. (1997); I93 = Inglis et al. (1993); K98 = Kay & Moran (1998); L01 = Lumsden et al. (2001); M90 = Miller & Goodrich (1990); M00 = Moran et al. (2000); M01 = Moran et al. (2001); R94 = Ruiz et al. (1994); T92 = Tran et al. (1992); T01 = Tran (2001); V01 = Veron & Veron (2001); Y96 = Young et al. (1996).

Table 2. Optical and hard X-ray data for Seyfert 2 galaxies

Name (1)	L_B^a (2)	$L_{H\beta}^b$ (3)	$L_{[OIII]}^c$ (4)	References (5)	$\log_{10} N_H$ (6)	$L_{2-10\text{keV}}^d$ (7)	References (8)	L_{Edd}^e (9)
Mrk 334		40.507	41.289	d88	20.643	< 43.102	p02	
NGC 34		39.056	42.772	d88				
IRAS 00198-7926		40.532	42.562	d92	> 24.0		r00	
Mrk 348	10.220	39.870	41.912	d88	23.041	43.011	b99	45.211
IRAS 00521-7054		40.940	42.743	d92				
NGC 424	10.160	40.350	41.462	m94	24.176		c00	
NGC 513	10.760	40.327	> 40.597	c94				45.730
NGC 591	10.349	39.972	41.954	d88				44.767
Mrk 573	10.455	40.452	42.001	d88				45.296
IRAS 01475-0740		39.784	41.689	d92				
NGC 788	10.569	39.735	40.990	w92				45.562
NGC 1068	10.787	40.675	42.645	m94	> 25.0		b99	45.383
NGC 1144	11.048	39.720	42.251	d88	20.699	< 43.281	p02	
Mrk 1066	10.379	40.205	42.175	m94	> 24.0		r99	44.972
IRAS 02581-1136		40.307	41.536	d92				
NGC 1241	10.827	< 38.529	42.472	d85				
NGC 1320		39.346	40.959	m94				45.176
NGC 1358	10.578	39.757	40.783	m94				45.996
NGC 1386	9.350	38.787	40.586	m94	> 25.0		b99	45.246
IRAS 03362-1642		39.878	41.529	d92				
IRAS 04103-2838								
IRAS 04210+0400		40.463	42.328	y96				
IRAS 04229-2528		38.861	41.903	y96				
IRAS 04259-0440		38.908	40.440	v95				
IRAS 04385-0828		< 39.102	40.117	d92				
NGC 1667	10.792	39.889	41.921	s95	> 24.0		b99	45.996
NGC 1685		40.081	42.582	c94				
IRAS 05189-2524		39.633	42.459	v95	22.690	43.325	b99	
Mrk 3	10.333	41.018	43.221	m94	24.041		b99	46.900
NGC 2273	10.125	40.314	41.449	l92	> 25.0		b99	45.313
ESO 428-G014	9.976	39.621	41.937	a91	> 25.0		m98	
Mrk 1210	9.958	40.472	42.195	t91	> 24.0		b99	
IRAS 08277-0242								
NGC 3081	10.092	39.908	41.331	m94	23.820	41.831	b99	
NGC 3079	10.439	37.145	40.427	h97	22.204	40.251	b99	45.293
NGC 3281	10.707	38.044	40.998	m94	23.903	42.798	b99	
IRAS 10340+0609								
NGC 3362	10.998	40.001	41.269	p96				44.697
UGC 6100	10.738	40.592	42.180	c94				45.784
IRAS 11058-1131		41.199	42.316	d92	> 24.0		r00	
NGC 3660		39.367	40.914	s95	20.255	41.800	tar	
NGC 3982	9.933	38.174	40.019	p96				43.878
NGC 4117	8.767							44.767
Was 49b		41.257	42.412	m92	22.799	42.970	a00	
NGC 4388	10.925	39.691	41.684	m94	23.623	42.744	b99	45.228
NGC 4501	11.212	38.873	39.804	h97				45.717
NGC 4507	10.588	40.430	41.569	m94	23.462	43.217	b99	
IC 3639	10.367	40.255	42.113	s95	> 25.0		r99	

Table 2—Continued

Name (1)	L_B^a (2)	$L_{H\beta}^b$ (3)	$L_{[OIII]}^c$ (4)	References (5)	$\log_{10} N_H$ (6)	$L_{2-10\text{keV}}^d$ (7)	References (8)	L_{Edd}^e (9)
NGC 4941	9.817	38.836	40.894	s89	23.653	40.820	b99	
MCG -3-34-64		41.261	42.322	d88	23.881	42.531	b99	
NGC 5135	10.705	40.153	42.311	m94	> 24.0		b99	
NGC 5194	10.472	37.919	40.168	h97	23.699	39.956	b99	44.911
NGC 5252	10.558	39.924	41.963	y96	22.633	43.064	b99	46.188
NGC 5256		40.295	41.825	d88	> 24.0		b99	
NGC 5283	9.717	39.596	40.836	d88				45.670
Mrk 1361	10.049	40.285	42.242	v95				
IRAS 13452-4155		40.475	42.098	y96				
NGC 5347	9.994	< 38.803	40.446	d92	> 24.0		r99	44.723
Mrk 463E	11.206	41.223	42.785	d88	23.204	42.646	b99	
Circinus	9.914	39.402	40.474	o94	> 24.0		b99	44.293
NGC 5506	10.056	39.585	41.608	s95	22.531	42.863	b99	
NGC 5643	10.421	39.063	41.225	m94	> 25.0		b99	
NGC 5695	10.380	39.383	40.501	d88				45.626
Mrk 477	10.488	41.626	43.543	d88	> 24.0		b99	
NGC 5728	10.644	39.946	42.083	s95				46.383
ESO 273-IG04		41.129	42.372	y96				
NGC 5929		39.545	40.926	m94	20.763	< 42.088	p02	45.262
NGC 5995		42.104	42.904	l01	21.934	43.426	l01	
IRAS 15480-0344		40.021	42.946	d92				
IRAS 17345+1124		41.961	42.956	t99				
NGC 6552	10.649		42.122	b99	23.778	42.461	b99	
IRAS 19254-7245		40.044	42.731	v02	> 24.0		r00	
NGC 6890	10.111	39.059	40.808	m94				
IRAS 20210+1121		41.454	43.124	y96	> 25.0		b99	
IRAS 20460+1925		41.113	42.838	y96	22.398	44.076	b99	
IC 5063	10.470	40.322	41.919	m94	23.380	42.849	b99	45.836
NGC 7130	10.677	40.547	42.477	s95	> 24.0		r00	
NGC 7172	10.279	38.733	39.767	v97	22.934	42.495	b99	
IRAS 22017+0319		41.225	42.497	d92	21.301	43.410	r00	
NGC 7212	10.657	40.941	42.636	m94	23.653	42.356	a00	45.517
MCG -3-58-7		< 40.484	41.692	d92				
IRAS 23060+0505		< 41.005	>43.916	d92	22.924	44.155		
NGC 7496	10.143	39.382	40.211	y96	22.699	< 41.652	p96	
IRAS 23128-5919								
IC 5298	10.489	39.927	42.078	v95				
NGC 7582	10.507	39.912	41.357	s95	23.079	42.144	b99	
NGC 7590	10.265	39.152	39.949	s95	< 20.964	40.798	b99	
NGC 7672	9.996							44.831
NGC 7674	10.914	40.808	42.495	d88	> 25.0		b99	45.620
NGC 7682	10.431	40.197	41.718	c94				45.296

^aBlue luminosities in units of $L_\odot$

^bObserved H_β luminosities in units of erg s^{-1} .

^cExtinction-corrected $[O III]\lambda 5007$ luminosities in units of erg s^{-1} .

^dAbsorption-corrected 2-10 keV luminosities for Compton-thin Seyfert 2 galaxies in units of erg s^{-1} .

^eEddington luminosities in units of erg s^{-1} .

Note. — REFERENCES: a91=Acker et al. 1991; a00=Awaki et al. 2000; b99=Bassani et al. 1999; c94=Cruz-Gonzalez et al. 1994; c00=Collinge & Brandt 2000; d85=Dahari 1985; d88=Dahari & De Robertis 1988; d92=de Grijp et al. 1992; h97=Ho et al. 1997; l92=Lonsdale 1992; l01=Levenson et al. 2001; lu01=Lumsden et al. 2001; m92=Moran et al. 1992; m94=Mulchaey et al. 1994; m98=Maiolino et al. 1998; o94=Oliva et al. 1994; p96=Polletta et al. 1996; p02=Almudena Prieto et al. 2002; r99=Risaliti et al. 1999; r00=Risaliti et al. 2000; s89=Storchi Bergmann & Pastoriza 1989; s95=Storchi-Bergmann et al. 1995; tar=tartarus.gsfc.nasa.gov; t91=Teleovich et al. 1991; t99=Tran et al. 1999; v95=Veilleux et al. 1995; v97=Veceli et al. 1997; v02=Vanzi et al. 2002; w92=Whittle 1992; y96=Young et al. 1996.

Table 3. Statistical results for Seyfert 2 galaxies

parameters	Sy2s with PBL	Sy2s without PBL	possibility ^a
$\log_{10}N_{\text{H}}$ (cm^{-2})	23.633 ± 0.201	23.562 ± 0.350	79% ^c
$\log_{10}L_{\text{B}}$ ($L_{\odot}$)	10.460 ± 0.068	10.340 ± 0.087	69% ^b
$\log_{10}L_{\text{FIR}}(L_{\odot})$	10.506 ± 0.106	10.301 ± 0.106	38% ^b
$\log_{10}(L_{\text{FIR}}/L_{\text{B}})$	-0.275 ± 0.069	-0.298 ± 0.078	21% ^b
f_{60}/f_{25}	2.447 ± 0.201	4.974 ± 0.375	0.002% ^b
$\log_{10}L_{25\mu\text{m}}$ ($L_{\odot}$)	10.565 ± 0.102	10.021 ± 0.111	0.005% ^b
$\log_{10}L_{1.49\text{GHz}}(\text{erg s}^{-1} \text{ Hz}^{-1})$	29.803 ± 0.136	29.109 ± 0.135	0.02% ^b
$\log_{10}L_{[\text{OIII}]}(\text{erg s}^{-1})$	42.147 ± 0.123	41.426 ± 0.136	0.07% ^c
$\log_{10}L_{2-10\text{keV}}(\text{erg s}^{-1})^d$	42.996 ± 0.138	41.220 ± 0.320	0.001% ^c

^aThe possibility from the same parent population.

^bTwo-sample Kolmogorov – Smirnov test.

^cSurvival data analysis program *asurv* Rev 1.2, Gehan’s Generalized Wilcoxon Test (Hypergeometric variance).

^dJust for Compton-thin Seyfert 2 galaxies.

Table 4. Common Sy2s observed in different samples

Samples ^a	Y96	M00	T01	L01
Y96	24(11) ^b	2(1)	9(9 ^c)	3(3)
M00		31(11)	13(5)	4(1)
T01			49(22)	16(8)
L01				22(8)

^aReferences for samples: Y96 = Young et al. 1996; M00 = Moran et al., 2000; T01 = Tran 2001; L01 = Lumsden et al. 2001

^bThe numbers indicate the numbers of common Seyfert 2 galaxies and the numbers in parentheses are numbers of Sy2s with PBL

^cOnly 6 Sy2s with PBL have been detected by Young et al. (1996).